Leading Causes of Death in Casey County, Kentucky, 2011

Cause of Death	Number of Deaths in Casey County[a] (% of all deaths)	Rate per 100,000 Total Population		
		Casey County[a]	Kentucky[b]	United States[b]
†Heart Disease	36 (19.1%)	185.6	205.6	173.7
†Malignant Neoplasms (Cancer)	49 (26.1%)	236.1	195.9	168.6
†Chronic Lower Respiratory	14 (7.4%)	71.8	63.1	42.7
Unintentional Injuries	12 (6.4%)	72.5	55.7	38.0
†Stroke	9 (4.8%)	50.5	43.1	37.9

† Leading causes of death known to be associated with secondhand smoke exposure.

[a] 2011 Vital Statistics Reports from the Kentucky Cabinet for Health and Family Services
Age-adjusted rates per 100,000 2011 population.
Estimates reflect only Kentucky residents who died in Kentucky

[b] Hoyert DL, Xu JQ. Deaths: Preliminary data for 2011. *National Vital Statistics Reports*, *(61)*6. Hyattsville, MD: National Center for Health Statistics, 2012.

How Secondhand Smoke is Connected with the Leading Causes of Death:

- Nonsmokers who are exposed to secondhand smoke at home or work increase their risks of developing *lung cancer* by 20–30%.[1]
- Nonsmokers who are exposed to secondhand smoke at home or work increase their *heart disease* risk by 25–30%.[1]

Public Health Solutions:

- After strong smoke-free laws go into effect, communities experience a 15% drop in heart attacks annually.[2]
- Smoke-free laws create healthy places for workers and patrons and reduce health care costs. Smoke-free laws are a known public health solution.[1]

[1]U.S. Department of Health and Human Services*. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General.* Rockville, MD: U.S. Department of Health and Human Services, 2006.
[2]Tan CE, Glantz SA.(2012). Association between smoke-free legislation and hospitalizations for cardiac, cerebrovascular, and respiratory diseases*. Circulation*, *126,* 2177-2183*.*

For more information, contact Kentucky Center for Smoke-free Policy
UK College of Nursing at 859-323-4587

April 2013