

The Kentucky Department for Public Health  
University of Kentucky College of Nursing  
**FIVCO District Health Department Tobacco Cessation Survey, 2002**

	<u>FIVCO</u>	<u>Kentucky (N=54)*</u>
1. Are you or other health department staff involved in providing tobacco cessation programs?	Yes	54 (100.0%) Yes
2. If "Yes," what programs do you provide?		
a. Fresh Start (American Cancer Society)	No	11 (20.4%) Yes
b. Freedom from Smoking (American Lung Association)	No	1 (1.9%) Yes
c. Cooper-Clayton Method to Stop Smoking	Yes	50 (92.6%) Yes
d. Make Yours a Fresh Start Family (American Cancer Society)	No	37 (68.5%) Yes
e. Quit and Win	No	31 (57.4%) Yes
3. If "Yes," how many tobacco cessation programs were offered within the last year? (Entire programs, not individual classes)	5	Mean = 4.7 ( <u>SD</u> = 3.5)
4. If "Yes," how many clients/patients participated in tobacco cessation programs within the last year?	40	Mean = 72.4 ( <u>SD</u> = 79.1)
5. If "Yes," how many clients/patients completed tobacco cessation programs within the last year?	25	Mean = 41.5 ( <u>SD</u> = 65.7)
6. If "Yes," what is the job title of health department staff providing tobacco cessation (Check all that apply)?		
a. Health Educator	Yes	34 (63.0%) Yes
b. Nurse	No	36 (66.7%) Yes
c. Social Worker	No	5 (9.3%) Yes
d. Physician	No	3 (5.6%) Yes
7. On average, how many requests for tobacco cessation does your health department receive per month?	0	Mean = 8.0 ( <u>SD</u> = 14.3)

The Kentucky Department for Public Health  
University of Kentucky College of Nursing  
**FIVCO District Health Department Tobacco Cessation Survey, 2002**

	<u>FIVCO</u>	<u>Kentucky (N=54)*</u>
8. According to our records, 1 person in your area participated in the Quit & Win Tobacco-Free Contest. To the best of your knowledge, about how many people involved in your cessation programs also participated in Quit and Win?	Don't Know	Mean = 4.4 ( <u>SD</u> = 11.3)
9. Within the past year, have any Cooper-Clayton Method to Stop Smoking programs been provided in your service area?	Yes	50 (92.6%) Yes
10. If "Yes", how many people in your service area are trained to provide Cooper-Clayton programs?	8	Mean = 5.1 ( <u>SD</u> = 8.2)
11. Within the past year, has your health department provided education programs on tobacco cessation for health care providers?	Yes	30 (57.7%) Yes
12. If "Yes," about how many programs did you provide?	2	Mean = 2.6 ( <u>SD</u> = 2.8)
13. Does your health department have a tobacco-user identification system (e.g. tobacco use as a vital sign, flagging the charts)?	Yes	45 (84.9%) Yes
14. If "Yes," is the tobacco-user identification system used for every patient at every clinic visit?	Yes	38 (82.6%) Yes
15. What kinds of educational methods are used for tobacco use cessation in your health department?		
a. Formal group classes	Yes	45 (83.3%) Yes
b. Individual counseling	Yes	52 (96.3%) Yes
c. Audio-visual materials	Yes	47 (87.0%) Yes
d. Printed materials	Yes	54 (100.0%) Yes
e. Posters	Yes	48 (88.9%) Yes
16. Does your clinic have self-help materials for tobacco use cessation available in every exam room?	Yes	38 (70.4%) Yes

The Kentucky Department for Public Health  
University of Kentucky College of Nursing  
**FIVCO District Health Department Tobacco Cessation Survey, 2002**

	<u>FIVCO</u>	<u>Kentucky (N=54)*</u>										
17. Are materials on secondhand smoke available in every exam room?	Yes	32 (62.8%) Yes										
18. Do your health care providers routinely ask patients about secondhand smoke exposure in the home?	Yes	46 (85.2%) Yes										
19. Do your health care providers routinely advise patients to ban smoking in the home?	Yes	44 (84.6%) Yes										
20. Do you health care providers routinely advise patients to ban smoking in their cars?	Yes	34 (69.4%) Yes										
21. Do your providers routinely refer tobacco users who want to quit to a counselor or to a program available in your community for tobacco cessation treatment?	Yes	52 (98.1%) Yes										
22. Does your employee health plan(s) at your health department reimburse for tobacco use cessation treatment? (including nicotine gum/patch and/or counseling)	No	20 (42.6%) Yes										
23. How frequently do your health care providers prescribe or recommend the following medications for tobacco use cessation?												
a. Nicotine Gum	3=	<table style="width: 100%; border: none;"> <tr><td>1= Never</td><td style="text-align: right;">13.2%</td></tr> <tr><td>2=</td><td style="text-align: right;">32.1%</td></tr> <tr><td>3=</td><td style="text-align: right;">37.7%</td></tr> <tr><td>4=</td><td style="text-align: right;">9.4%</td></tr> <tr><td>5= Very Often</td><td style="text-align: right;">7.5%</td></tr> </table>	1= Never	13.2%	2=	32.1%	3=	37.7%	4=	9.4%	5= Very Often	7.5%
1= Never	13.2%											
2=	32.1%											
3=	37.7%											
4=	9.4%											
5= Very Often	7.5%											
b. Nicotine Patch	4=	<table style="width: 100%; border: none;"> <tr><td>1= Never</td><td style="text-align: right;">7.5%</td></tr> <tr><td>2=</td><td style="text-align: right;">26.4%</td></tr> <tr><td>3=</td><td style="text-align: right;">34.0%</td></tr> <tr><td>4=</td><td style="text-align: right;">13.2%</td></tr> <tr><td>5= Very Often</td><td style="text-align: right;">18.9%</td></tr> </table>	1= Never	7.5%	2=	26.4%	3=	34.0%	4=	13.2%	5= Very Often	18.9%
1= Never	7.5%											
2=	26.4%											
3=	34.0%											
4=	13.2%											
5= Very Often	18.9%											
c. Nicotine Nasal Spray	3=	<table style="width: 100%; border: none;"> <tr><td>1= Never</td><td style="text-align: right;">62.3%</td></tr> <tr><td>2=</td><td style="text-align: right;">20.8%</td></tr> <tr><td>3=</td><td style="text-align: right;">17.0%</td></tr> <tr><td>4=</td><td style="text-align: right;">0.0%</td></tr> <tr><td>5= Very Often</td><td style="text-align: right;">0.0%</td></tr> </table>	1= Never	62.3%	2=	20.8%	3=	17.0%	4=	0.0%	5= Very Often	0.0%
1= Never	62.3%											
2=	20.8%											
3=	17.0%											
4=	0.0%											
5= Very Often	0.0%											

The Kentucky Department for Public Health  
 University of Kentucky College of Nursing  
**FIVCO District Health Department Tobacco Cessation Survey, 2002**

	<u>FIVCO</u>	<u>Kentucky (N=54)*</u>	
d. Nicotine Inhaler		1= Never	60.4%
	2=	2=	28.3%
		3=	11.3%
		4=	0.0%
		5= Very Often	0.0%
e. Bupropion (Wellbutrin, Zyban)		1= Never	35.8%
		2=	26.4%
		3=	22.6%
		4=	11.3%
	5= Very Often	5= Very Often	3.8%
f. Clonidine (suppresses opioid withdrawal symptoms; typically used to treat high blood pressure)		1= Never	82.7%
	2=	2=	9.6%
		3=	7.7%
		4=	0.0%
		5= Very Often	0.0%
g. Nortriptyline (antidepressant)		1= Never	80.4%
	2=	2=	9.8%
		3=	7.8%
		4=	0.0%
		5= Very Often	2.0%
24. We have enough staff to provide tobacco cessation programs.	1= StronglyDisagree	1= StronglyDisagree	14.8%
		2= Disagree	31.5%
		3= Agree	37.0%
		4= StronglyAgree	16.7%
25. Our staff are not trained to provide tobacco cessation.	2= Disagree	1= StronglyDisagree	46.3%
		2= Disagree	38.9%
		3= Agree	13.0%
		4= StronglyAgree	1.9%
26. We are too busy to provide tobacco cessation.		1= StronglyDisagree	42.6%
		2= Disagree	44.4%
		3= Agree	11.1%
	4= StronglyAgree	4= StronglyAgree	1.9%

\* Participation rate = 98.2%

The Kentucky Department for Public Health  
 University of Kentucky College of Nursing  
**FIVCO District Health Department Tobacco Cessation Survey, 2002**

	<u>FIVCO</u>	<u>Kentucky (N=54)*</u>	
27 We have few requests for tobacco cessation programs.		1= StronglyDisagree	24.1%
		2= Disagree	29.6%
	3= Agree	3= Agree	37.0%
		4= StronglyAgree	9.3%
28. Tobacco cessation programs are too expensive.	1= StronglyDisagree	1= StronglyDisagree	35.2%
		2= Disagree	44.4%
		3= Agree	18.5%
		4= StronglyAgree	1.9%
29. Tobacco cessation programs are effective.		1= StronglyDisagree	0.0%
		2= Disagree	1.9%
	3= Agree	3= Agree	67.3%
		4= StronglyAgree	30.8%

**For more information, contact:**  
**Dr. Ellen Hahn, DNS, RN**  
**Associate Professor**  
**University of Kentucky**  
**College of Nursing**  
**(859) 257-2358**  
[ejhahn00@pop.uky.edu](mailto:ejhahn00@pop.uky.edu)

\* Participation rate = 98.2%