

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

	<u>Northern Kentucky</u>	<u>Kentucky (N=55)</u>
1. Are you or other health department staff involved in providing tobacco cessation programs?	Yes	40 (72.7%) Yes
2. If "Yes", what programs do you provide?		
a. Fresh Start (American Cancer Society)	No	7 (17.5%)
b. Freedom from Smoking (American Lung Association)	No	0
c. Cooper-Clayton Method to Stop Smoking	Yes	35 (87.5%)
d. Make Yours a Fresh Start Family (American Cancer Society)	No	8 (20.0%)
3. If "Yes", how many tobacco cessation programs were offered within the last year? (Entire programs, not individual classes)	4	Mean = 3.8 (<u>SD</u> = 4.0)
4. If "Yes", how many clients/patients participated in tobacco cessation programs within the last year?	150	Mean = 34.0 (<u>SD</u> = 44.5)
5. If "Yes", how many clients/patients completed tobacco cessation programs within the last year?	60	Mean = 15.8 (<u>SD</u> = 24.2) (45.3% Completion Rate)
6. If "Yes", what is the job title of health department staff providing tobacco cessation (Check all that apply)?		
a. Health Educator	Yes	26 (65.0%)
b. Nurse	No	15 (37.5%)
c. Social Worker	No	3 (12.5%)
7. On average, how many requests for tobacco cessation does your health department receive per month?	10	Mean = 7.4 (<u>SD</u> = 14.6)

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

	<u>Northern Kentucky</u>	<u>Kentucky (N=55)</u>
8. Within the past year, have any Cooper-Clayton Method to Stop Smoking programs been provided in your service area?	Yes	29 (53.7%) Yes
9. If “Yes”, how many people in your service area are trained to provide Cooper-Clayton programs?	16	Mean = 4.9 (<u>SD</u> = 5.3)
10. Within the past year, has your health department provided education programs on tobacco cessation for health care providers?	No	15 (27.3%) Yes
11. If “Yes”, about how many programs did you provide?	--	Mean = 3.1 (<u>SD</u> = 2.4)
12. Does your health department have a tobacco-user identification system (e.g. tobacco use as a vital sign, flagging the charts)?	Yes	36 (65.5%) Yes
13. If “Yes”, is the tobacco-user identification system used for every patient at every clinic visit?	Yes	32 (88.9%) Yes
14. What kinds of educational methods are used for tobacco use cessation in your health department?		
a. Formal group classes	Yes	37 (67.3%)
b. Individual counseling	Yes	51 (92.7%)
c. Audio-visual materials	No	36 (65.5%)
d. Printed materials	No	54 (98.2%)
e. Posters	No	48 (87.3%)
15. Does your clinic have self-help materials for tobacco use cessation available in every exam room?	No	32 (58.2%) Yes

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

	<u>Northern Kentucky</u>	<u>Kentucky (N=55)</u>
16. Are materials on secondhand smoke available in every exam room?	No	27 (49.0%) Yes
17. Do your health care providers routinely ask patients about secondhand smoke exposure in the home?	Yes	41 (75.9%) Yes
18. Do your health care providers routinely advise patients to ban smoking in the home?	Yes	39 (73.6%) Yes
19. Do you health care providers routinely advise patients to ban smoking in their cars?	Yes	34 (64.2%) Yes
20. Do your providers routinely refer tobacco users who want to quit to a counselor or to a program available in you community for tobacco use cessation treatment?	Yes	47 (85.5%) Yes
21. Does your employee health plan(s) at your health department reimburse for tobacco use cessation treatment? (including nicotine gum/patch and/or counseling)	Yes	15 (27.8%) Yes
22. How frequently do your health care providers prescribe or recommend the following medications for tobacco use cessation?		

a. Nicotine Gum

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Never	40.0%
	2 =	32.7%
3 =	3 =	18.2%
	4 =	7.3%
	5 = Very Often	5.5%

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

b. Nicotine Patch

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Never	38.2%
	2 =	23.6%
3 =	3 =	18.2%
	4 =	10.9%
	5 = Very Often	9.1%

c. Nicotine Nasal Spray

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Never	1 = Never	80.0%
	2 =	18.2%
	3 =	1.8%
	4 =	0.0%
	5 = Very Often	0.0%

d. Nicotine Inhaler

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Never	1 = Never	76.4%
	2 =	21.8%
	3 =	1.8%
	4 =	0.0%
	5 = Very Often	0.0%

e. Bupropion (Wellbutrin, Zyban)

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Never	1 = Never	42.6%
	2 =	27.8%
	3 =	14.8%
	4 =	3.7%
	5 = Very Often	11.1%

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

f. Clonidine (suppresses opioid withdrawal symptoms; typically used to treat high blood pressure)

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Never	1 = Never	92.6%
	2 =	7.4%
	3 =	0.0%
	4 =	0.0%
	5 = Very Often	0.0%

g. Nortriptyline (antidepressant)

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Never	1 = Never	94.4%
	2 =	3.7%
	3 =	1.9%
	4 =	0.0%
	5 = Very Often	0.0%

23. We have enough staff to provide tobacco cessation programs.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Strongly Disagree	9.1%
2 = Disagree	2 = Disagree	36.4%
	3 = Agree	40.0%
	4 = Strongly Agree	14.5%

24. Our staff are not trained to provide tobacco cessation.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Strongly Disagree	1 = Strongly Disagree	12.7%
	2 = Disagree	61.8%
	3 = Agree	18.2%
	4 = Strongly Agree	7.3%

The Kentucky Department for Public Health
 University of Kentucky College of Nursing
Northern Kentucky Independent District Health Department Tobacco Cessation Survey, 2001

25. We are too busy to provide tobacco cessation.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Strongly Disagree	21.8%
2 = Disagree	2 = Disagree	72.7%
	3 = Agree	5.5%
	4 = Strongly Agree	0.0%

26. We have few requests for tobacco cessation programs.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
1 = Strongly Disagree	1 = Strongly Disagree	10.9%
	2 = Disagree	41.8%
	3 = Agree	36.4%
	4 = Strongly Agree	10.9%

27. Tobacco cessation programs are too expensive.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Strongly Disagree	9.1%
	2 = Disagree	67.3%
3 = Agree	3 = Agree	20.0%
	4 = Strongly Agree	3.6%

28. Tobacco cessation programs are effective.

<u>Northern Kentucky</u>	<u>Kentucky</u>	
	1 = Strongly Disagree	0.0%
	2 = Disagree	10.9%
	3 = Agree	67.3%
4 = Strongly Agree	4 = Strongly Agree	21.8%

**For more information,
 contact:
 Dr. Ellen Hahn
 University of Kentucky
 College of Nursing
 (859) 257-2358
eihahn00@pop.uky.edu**